

IFT-UAM-CSIC-02-03

hep-th/0301032

Chiral four-dimensional string compactifications with intersecting D-branes

Angel M. Uranga ¹

I.M.A.F.F. and Instituto de Física Teórica C-XVI

Universidad Autónoma de Madrid, 28049 Madrid, Spain

We review the construction of chiral four-dimensional compactifications of type IIA string theory with intersecting D6-branes. Such models lead to four-dimensional theories with non-abelian gauge interactions and charged chiral fermions. We discuss the application of these techniques to building of models with spectrum as close as possible to the Standard Model, and review their main phenomenological properties. We also emphasize the advantages/disadvantages of carrying out this idea using supersymmetric of non-supersymmetric models.

¹Angel.Uranga@uam.es

1 Introduction

String theory has the remarkable property that it provides a description of gauge and gravitational interactions in a unified framework consistently at the quantum level. It is this general feature (beyond other beautiful properties of *particular* string models) that makes this theory interesting as a possible candidate to unify our description of the different particles and interactions in Nature.

Now if string theory is indeed realized in Nature, it should be able to lead not just to ‘gauge interactions’ in general, but rather to gauge sectors as rich and intricate as the gauge theory we know as the Standard Model of Particle Physics. This is described by a gauge group

$$SU(3)_c \times SU(2)_w \times U(1)_Y \quad (1)$$

a set of charged chiral (left-handed) fermions in three copies with identical gauge quantum numbers, namely

$$3 \times [(3, 2)_{1/6} + (\bar{3}, 1)_{1/3} + (\bar{3}, 1)_{-2/3} + (1, 2)_{-1/2} + (1, 1)_1 + (1, 1)_0] \quad (2)$$

(where we have also included the right-handed neutrinos), and a scalar particle, the Higgs multiplet, with gauge quantum numbers

$$(1, 2)_{-1/2} \quad (3)$$

These particles have also very specific interactions. They correspond to all possible terms consistent with the gauge symmetries, and with some ‘accidental’ global symmetries, like baryon or lepton number.

This quantum field theory has certain characteristic and very interesting rough features, for instance the very existence of non-abelian gauge interactions, the presence of charged chiral fermions, the spontaneous breaking of gauge symmetries, the replication of fermion families, etc. On the other hand, once this basic structure is fixed, the model contains a large number of more detailed features, for instance the particular values of gauge couplings or the pattern of yukawa interaction strengths, etc.

All these features are simply unexplained by the Standard Model itself, where they are external inputs. It is a natural question to ask whether it is possible to explain or reproduce

them in a microscopic underlying theory, like string theory. In a sense to be qualified below, this is the purpose of the branch of string theory known as String Phenomenology.

Clearly, even if string theory is correct, it is not realistic to hope to construct explicitly ‘the’ string theory of the world, in the following sense. As is well known, the simplest string theories propagate in spacetimes of ten dimensions (eleven for M-theory), and usually have a high degree of supersymmetry. In the process of describing models reducing at low energies to four dimensions and low or no supersymmetry, there is an enormous arbitrariness in the choice of the background configuration, namely the compactification data, etc. We do not have a good enough understanding of whether any of these is in any sense preferred over the rest (either dynamical, cosmological or anthropically). Moreover, most of the regimes of string theory are not accessible to our computational tools (which involve perturbation theory in spacetime string coupling, and in α' corrections around exactly solvable conformal field theory backgrounds). These considerations imply that it is extremely unlikely that we find the ‘correct’ string theory by trial and error.

From this viewpoint, the purpose of String Phenomenology (for this talk) is more modest, but still non-trivial and hopefully achievable:

I. We should describe (and hopefully classify) different *setups* in string theory, which lead to the same kind of physics as the Standard Model (at the rough level, namely leading to non-abelian gauge interactions, replicated chiral fermions, etc).

II. Within each setup, we should construct explicit examples with low energy physics as close as possible to the Standard Model. From these particular examples we should extract generic, robust, features of their low-energy phenomenology, which can therefore be considered as natural predictions of the setup.

Once this program is fulfilled to a satisfactory level (and we are still far from understanding diverse issues, e.g. related to the absence of exact supersymmetry at low-energy), future experimental data should be able to start cornering what kind of string theory may be underlying our world.

Prototypical example: Compactification of heterotic string

At this point it will be useful to provide the most notorious example of what we mean by a ‘setup’, and what lessons can be drawn from it. The most familiar string theory setup reproducing the rough features of the Standard Model at low energies is compactification of the heterotic string theories (although we center on perturbative heterotic models, similar features are also valid for compactifications of Horava-Witten theory, or in non-perturbative




Figure 1: Picture of heterotic string compactification.

heterotic vacua with five-branes).

I. Construction

The heterotic string leads to a highly supersymmetric massless sector containing gravitational and gauge interactions (with gauge group G being $E_8 \times E_8$ or $SO(32)$), propagating in ten-dimensional spacetime.

To obtain four-dimensional physics, and to reduce the amount of supersymmetry, we choose the background spacetime to be $\mathbf{M}_4 \times \mathbf{X}_6$, with $\mathbf{X}_6$ a compact six-dimensional manifold (usually chosen Calabi-Yau to preserve 4d $\mathcal{N} = 1$ supersymmetry). Moreover, to reduce the 4d gauge group, we turn on a non-trivial background for the internal components of gauge fields in a subgroup $H \subset G$. The 4d gauge group is given by the commutant of H in G , namely the elements of G commuting with H (for instance, for $G = E_8 \times E_8$, the choice $H = SU(3)$ leads to a 4d gauge group $E_6 \times E_8$; while $H = SU(3) \times \mathbf{Z}_2$ can lead to a 4d gauge group $SU(3) \times SU(2) \times U(1)$). Fig. ?? shows configurations of this kind.

Finally, these backgrounds ensure that the low-energy theory naturally contains 4d chiral fermions charged under the gauge group, replicated a number of times determined by the topology of the compactification space $\mathbf{X}_6$ and of the gauge background (for instance, for the choice of internal gauge background known as standard embedding, with $H = SU(3)$, the number of families, i.e. 27's of E_6 is given by the Euler characteristic $\chi(\mathbf{X}_6)$ of $\mathbf{X}_6$).

II. Generic features/natural predictions of the setup

Explicit heterotic models of this kind, with spectrum very close to that of the Standard Model, have been constructed using different techniques (Calabi-Yau compactifications with non-standard embedding, orbifolds, fermionic constructions, etc). This allows to extract certain generic features of this setup.

- A very important one is the value of the string scale in this setup, which follows from analyzing the strength of gravitational and gauge interactions, as we quickly review.

The 10d gravitational and gauge interactions have the structure

$$\int d^{10}x \frac{M_s^8}{g_s^2} R_{10d} \quad ; \quad \int d^{10}x \frac{M_s^6}{g_s^2} F_{10d}^2 \quad (4)$$

where M_s , g_s are the string scale and coupling constant, and R_{10d} , F_{10d} are the 10d Einstein and Yang-Mills terms. Upon Kaluza-Klein compactification on $\mathbf{X}_6$, these interactions reduce to 4d and pick up a factor of the volume V_6 of $\mathbf{X}_6$

$$\int d^4x \frac{M_s^8 V_6}{g_s^2} R_{10d} \quad ; \quad \int d^4x \frac{M_s^6 V_6}{g_s^2} F_{10d}^2 \quad (5)$$

From this we may express the experimental 4d Planck scale and gauge coupling in terms of the microscopic parameters of the string theory configuration

$$M_P^2 = \frac{M_s^8 V_6}{g_s^2} \simeq 10^{19} \text{ GeV}^2 \quad ; \quad \frac{1}{g_{YM}^2} = \frac{M_s^6 V_6}{g_s^2} \simeq \mathcal{O}(1) \quad (6)$$

From these we obtain the relation

$$M_s = g_{YM} M_P \simeq 10^{18} \text{ GeV} \quad (7)$$

which implies that the string scale is necessarily very large in this kind of constructions.

- In order to protect the hierarchy between this fundamental scale and the weak scale $M_w \simeq 10^2 \text{ GeV}$, the requirement of $\mathcal{N} = 1$ susy at intermediate scales is quite essential.

- The large fundamental scale helps in protecting the proton against too fast decay, by suppressing higher dimension operators violating baryon number.

- Gauge coupling constants unify at the string scale, and lead to values in relatively good agreement with measured low-energy couplings.

These and many other interesting features make this setup an extremely appealing picture of our world. It is important however to mention that our understanding of it is not fully satisfactory, and that difficult open questions remain, like the breaking of supersymmetry, or the mechanisms to stabilize moduli. Hopefully further ingredients in the construction will help to improve the situation in the future.

Our purpose in the present lecture is to show that there exist other setups in string theory, leading to constructions with features close to that of the Standard Model, and as satisfactory as (or even more than) the heterotic setup. In this lecture we will center on

compactifications of type IIA theory with D6-branes wrapped on intersecting 3-cycles, extensively studied in the last two years [1, 2, 3, 4, 5, 7, 8, 9]¹. The discussion is organized as follows: In Section 2 we briefly review D-branes in string theory, and we introduce intersecting D6-branes as a natural way to obtain charged 4d chiral fermions. In Section 3 we describe compactifications of type IIA theory with intersecting D6-branes, give rules to obtain their spectrum, and provide explicit examples for toroidal backgrounds. We briefly discuss the generic phenomenological features in this kind of construction, i.e. the natural predictions of the setup. In Section 4 we discuss the advantages/disadvantages of constructing supersymmetric vs non-supersymmetric models, and the different issues that arise in the latter case, and briefly describe some advanced models in each approach. Section 5 contains some final comments.

2 Intersecting D-branes

2.1 D-branes in string theory

New knowledge on string theory beyond perturbation theory has led to the introduction of new objects in string theory, D-branes, which provide a brand-new way of realizing non-abelian gauge symmetries in string theory. Therefore it is natural to explore the possibility of using them in searching for new setups with potential phenomenological application. We now review the main properties of D-branes from this biased viewpoint.

Type II string theories contains certain ‘soliton-like’ states in their spectrum, with $p + 1$ extended dimensions, the p -branes. They were originally found as solutions of the low-energy supergravity equations of motion. Subsequently, it was realized [13] that certain of these objects (known as Dp -branes) admit a fully stringy description, as $(p + 1)$ -dimensional subspaces on which open strings can end. Notice that these open strings are not present in the vacuum of the underlying string theory, but rather represent the fluctuations of the theory around the topological defect background. Namely, the closed string sector still describes the dynamics of the vacuum (gravitational interactions, etc), while open strings rather describe the dynamics of the object. The situation is shown in figure 2

The spectrum of fluctuations of the theory in the presence of the Dp -brane is obtained by quantizing closed strings and open strings ending on the Dp -brane. Since the open string

¹See [11, 10, 12] for earlier related work.




Figure 2: Fluctuations of string theory around a Dp -brane configuration are describe as open strings ending on its volume.

endpoints are fixed on the D-brane, the massless modes in the latter sector yield fields propagating on the $(p + 1)$ -dimensional D-brane world-volume W_{p+1} . For a single type II Dp -brane in flat 10d space, such massless modes correspond to a $U(1)$ gauge boson, $9 - p$ real scalars and some fermions. The scalars (resp. fermions) can be regarded as Goldstone bosons (resp. Goldstinos) of the translational symmetries (resp. supersymmetries) of the vacuum broken by the presence of the D-brane. The open string sector fills out a $U(1)$ vector multiplet with respect to the 16 supersymmetries unbroken by the D-brane.

Dp -branes are charged under the corresponding RR $(p + 1)$ -form C_{p+1} of type II string theory, via the minimal coupling $\int_{W_{p+1}} C_{p+1}$. Since flat Dp -branes in flat space preserve $1/2$ of the 32 supercharges of the type II vacuum, such D-branes are BPS states, and their RR charge is related to their tension. This implies that there is no net force among parallel branes (roughly, gravitational attraction cancels against ‘Coulomb’ repulsion due to their RR charge), so they can be superposed. The open string spectrum in a configuration of n coincident Dp -branes consists of n^2 sectors, corresponding to the $n \times n$ possible ways of choosing the D-brane on which the string starts and ends. This multiplicity renders interactions between open strings non-abelian, and the complete massless open string spectrum is given by $U(n)$ gauge bosons, $9 - p$ adjoint scalars and adjoint fermions, filling out a $U(n)$ vector multiplet with respect to the 16 unbroken supersymmetries. The structure of gauge bosons for $n = 3$ is shown in figure 3. As announced, D-branes provide a brand-new realization of non-abelian gauge symmetry in string theory. Indeed, the low-energy effective action for the massless open string modes is Dirac-Born-Infeld action, which at low energies (lowest derivative order) reduces to the Yang-Mills action.




Figure 3: Non-abelian gauge bosons in a configuration of coincident D-branes.




Figure 4: Picture of D-brane worlds.

2.2 Brane-worlds

Hence D-branes allow for gauge sectors localized on subspaces of spacetime, in a consistent and microscopically well-defined framework. This is a string theory realization of the brane-world idea [14]. Namely we have a configuration where gravity propagates in 10d (with 4d gravity eventually reproduced at low energies via standard compactification on some internal space $\mathbf{X}_6$); on the other hand, gauge interactions (hopefully, in a elaborated enough construction, with some gauge sector similar to the Standard Model) propagate just in the volume of a lower-dimensional subspace of spacetime. Starting with $(p+1)$ -dimensional gauge interactions on a Dp -brane, 4d gauge interactions can be obtained by considering the Dp -brane volume to be of the form $\mathbf{M}_4 \times \Sigma_{p-3}$, with Σ_{p-3} a $(p-3)$ closed subspace i.e. a $(p-3)$ -cycle, of $\mathbf{X}_6$. This is shown in picture 4.

Since gauge and gravitational interactions arise in different string sectors (open vs closed) and propagate on different dimensions, the analysis of couplings and scales differs from that in heterotic. Before compactification, gravitational and gauge interactions are described by

an effective action

$$\int d^{10}x \frac{M_s^8}{g_s^2} R_{10d} + \int d^{p+1}x \frac{M_s^{p-3}}{g_s} F_{(p+1)d}^2 \quad (8)$$

where the powers of g_s follow from the Euler characteristic of the worldsheet which produces interactions for gravitons (sphere) and for gauge bosons (disk).

Upon compactification, the 4d action picks up volume factors and reads

$$\int d^4x \frac{M_s^8 V_6}{g_s^2} R_{4d} + \int d^4x \frac{M_s^{p-3} V_\Sigma}{g_s} F_{4d}^2 \quad (9)$$

This allows to read off the 4d Planck mass and gauge coupling, which are experimentally measured.

$$\begin{aligned} M_P^2 &= \frac{M_s^8 V_{X_6}}{g_s^2} \simeq 10^{19} \text{ GeV} \\ 1/g_{YM}^2 &= \frac{M_s^{p-3} V_\Sigma}{g_s} \simeq 0.1 \end{aligned} \quad (10)$$

If the geometry is factorizable, we can split $V_{X_6} = V_\Sigma V_\perp$, with $V_\perp$ the transverse volume, and obtain

$$M_P^2 g_{YM}^2 = \frac{M_s^{11-p} V_\perp}{g_s} \quad (11)$$

This shows that it is possible to generate a large Planck mass in 4d with a low string scale, by simply increasing the volume transverse to the brane. In particular, it has been proposed to lower the string scale down to the TeV scale to avoid a hierarchy with the weak scale. The hierarchy problem is recast in geometric terms, namely the stabilization of the compactification size in very large volumes (the extreme case being around one millimeter for two large dimensions). Notice however that a low string scale is not compulsory in models with some solution to the hierarchy problem, e.g. supersymmetric models.

2.3 D-branes and chirality

We have remarked that D-branes in flat space preserve a lot of supersymmetry, namely 16 supercharges, which amounts to $\mathcal{N} = 4$ in 4d. This is too much to allow for a chiral open string spectrum. Indeed it is quite non-trivial to construct configurations of D-branes with chiral open string sector. One can heuristically show that *isolated* D-branes with a *smooth*




Figure 5: Isolated D-branes at a smooth point in transverse space feel a locally trivial geometry and lead to non-chiral open string spectra.

transverse space automatically lead to non-chiral spectra ², as follows (see figure 5): The massless open string sector in a D-brane configuration is only sensitive to the local geometry around the D-branes. If they are isolated and sitting at a smooth point, the local structure is that of D-branes in flat space, which lead to non-chiral spectra.

There are two well-studied kinds of D-brane configurations which lead to chiral open string spectra, obtained by relaxing either of the above emphasized keywords.

a) D-branes sitting at *singular* (rather than smooth) points in transverse space can lead to chiral open string spectra. The prototypical example is given by stacks of D3-branes sitting at the singular point of orbifolds of flat space, e.g. orbifold singularities $\mathbf{C}^3/\mathbf{Z}_N$, as studied in [15]. Such configurations have been used for phenomenological model building e.g. in [16].

b) Sets of intersecting D-branes (which are hence not isolated) can also lead to chiral fermions in the sector of open strings stretched between different kinds of D-brane [17]. The chiral fermions are localized at the intersection of the brane volumes, in order to minimize its stretching. In the present lecture we will center on this kind of configuration.

2.4 Intersecting D-branes

The basic configuration of intersecting D-branes leading to chiral 4d fermions at their intersection is two stacks of D6-branes in flat 10d intersecting over a 4d subspace of their volumes. Figures 6 a, b provide two pictorial representations of the configurations.

²We are oversimplifying, since it may be possible to obtain chiral 4d fermions by turning on topologically non-trivial world-volume gauge backgrounds on Σ_{p-3} . We will not consider this possibility in the present




Figure 6: Two picture of D6-branes intersecting over a 4d subspace of their volumes.

Chirality of the sector of open string stretching between the two D6-branes is consistent with the fact that any continuous motion of the branes (preserving the gauge symmetry) maintains the existence of an intersection; this corresponds to the fact that chiral particles at the intersection do not become massive upon deforming their effective action in a continuous fashion.

The open string spectrum in a configuration of two stacks of (n_1 resp. n_2 coincident) D6-branes in flat 10d intersecting over a 4d subspace of their volumes consists of three open string sectors:

$\mathbf{6}_1\mathbf{6}_1$ Strings stretching between D6₁-branes provide $U(n_1)$ gauge bosons, three real adjoint scalars and fermion superpartners, propagating over the 7d world-volume of the D6₁-branes.

$\mathbf{6}_2\mathbf{6}_2$ Similarly, strings stretching between D6₂-branes provide $U(n_2)$ gauge bosons, three real adjoint scalars and fermion superpartners, propagating over the D6₂-brane 7d world-volume.

$\mathbf{6}_1\mathbf{6}_2 + \mathbf{6}_2\mathbf{6}_1$ Strings stretching between both kinds of D6-brane lead to a 4d chiral fermion, transforming in the representation $(n_1, \bar{n}_2)$ of $U(n_1) \times U(n_2)$, and localized at the intersection ³.

In addition, the latter sector leads to scalar fields in the same bi-fundamental representation, and whose masses depend on the local geometry of the intersection. The lightest

discussion.

³The chirality of the fermion is encoded in the orientation defined by the intersection; this will be implicitly taken into account in our discussion.




Figure 7: Compactification with intersecting D6-branes wrapped on 3-cycles.

complex scalars have masses

$$\begin{aligned} \alpha' M^2/2 = & \frac{1}{2}(\theta_1 + \theta_2 - \theta_3) \quad ; \quad \frac{1}{2}(\theta_1 - \theta_2 + \theta_3) \\ & \frac{1}{2}(-\theta_1 + \theta_2 + \theta_3) \quad ; \quad 1 - \frac{1}{2}(\theta_1 + \theta_2 + \theta_3) \end{aligned} \quad (12)$$

where the θ_i are $1/\pi$ times the angles defined in Fig 6b, and taken between -1 and 1 .

In the generic case, there is no supersymmetry invariant under the two stacks of branes, and the open string sector at the intersection is non-supersymmetric. However, if $\theta_1 \pm \theta_2 \pm \theta_3 = 0$ for some choice of signs, one of the scalars becomes massless, reflecting that the configuration is $\mathcal{N} = 1$ supersymmetric. $\mathcal{N} = 2$ supersymmetry arises if e.g. $\theta_3 = 0$ and $\theta_1 \pm \theta_2 = 0$, while $\mathcal{N} = 4$ arises only for parallel stacks $\theta_i = 0$.

3 Four-dimensional models

Once we have succeeded in describing configurations of D-branes leading to charged chiral fermions, in this section we employ them in building models with 4d gravity and gauge interactions. Although intersecting D6-branes provide 4d chiral fermions already in flat 10d space, gauge interactions remain 7d and gravity interactions remain 10d unless we consider compactification of spacetime.

The general kind of configurations we are to consider (see figure 3) is type IIA string theory on a spacetime of the form $\mathbf{M}_4 \times \mathbf{X}_6$ with compact $\mathbf{X}_6$, and with stacks of N_a D6_a-branes with volumes of the form $M_4 \times \Pi_a$, with $\Pi_a \subset \mathbf{X}_6$ a 3-cycles. It is important to realize that generically 3-cycles in a 6d compact space intersect at points, so the corresponding wrapped D6-branes will intersect at M_4 subspaces of their volumes. Hence, compactification




Figure 8: Examples of intersecting 3-cycles in $\mathbf{T}^6$.

reduces the 10d and 7d gravitational and gauge interactions to 4d, and intersections lead to charged 4d chiral fermions. Also, generically two 3-cycles in a 6d space intersect several times, therefore leading to a replicated sector of opens strings at intersections. This is a natural mechanism to explain/reproduce the appearance of replicated families of chiral fermions in Nature!

3.1 Toroidal models

In this section we mainly follow [2], see also [1]. To keep the configurations simple, we consider $\mathbf{X}_6$ to be a six-torus factorized as $\mathbf{T}^6 = \mathbf{T}^2 \times \mathbf{T}^2 \times \mathbf{T}^2$. Also for simplicity we take the 3-cycles Π_a to be given by a factorized product of 1-cycles in each of the 2-tori. For a 3-cycle Π_a , the 1-cycle in the i^{th} 2-torus will be labeled by the numbers (n_a^i, m_a^i) it wraps along the horizontal and vertical directions, see figure 8 for examples.

The intersection number is given by the product of the number of intersections in each 2-torus, and reads

$$I_{ab} = (n_a^1 m_b^1 - m_a^1 n_b^1) \times (n_a^2 m_b^2 - m_a^2 n_b^2) \times (n_a^3 m_b^3 - m_a^3 n_b^3) \quad (13)$$

It is useful to introduce the 3-homology class $[\Pi_a]$ of the 3-cycle Π_a , which can be thought of as a vector of RR charges of the corresponding D6-brane. The 1-homology class of an (n, m) 1-cycle in a 2-torus is $n[a] + m[b]$, with $[a], [b]$ the basic homology cycles in $\mathbf{T}^2$. For a 3-cycles with wrapping numbers (n_a^i, m_a^i) we have

$$[\Pi_a] = \otimes_{i=1}^3 (n_a^i [a_i] + m_a^i [b_i]) \quad (14)$$

The intersection number (13) is intersection number in homology, denoted $I_{ab} = [\Pi_a] \cdot [\Pi_b]$.

This is easily shown using $[a_i] \cdot [b_j] = \delta_{ij}$ and linearity and antisymmetry of the intersection pairing.

3.1.1 Spectrum

With the basic data defining the configuration, namely $\mathbf{N}_a$ D6_a-branes wrapped on 3-cycles $[\mathbf{\Pi}_a]$, with wrapping numbers $(\mathbf{n}_a^i, \mathbf{m}_a^i)$ on each $\mathbf{T}^2$ and intersection numbers $\mathbf{I}_{ab}$, we can compute the spectrum of the model.

The closed string sector produces 4d $\mathcal{N} = 8$ supergravity. There exist different open string sectors:

$\mathbf{6}_a \mathbf{6}_a$ String stretched among D6-branes in the a^{th} stack produce 4d $U(N_a)$ gauge bosons, 6 real adjoint scalars and 4 adjoint Majorana fermions, filling out a vector multiplet of the 4d $\mathcal{N} = 4$ supersymmetry preserved by the corresponding brane.

$\mathbf{6}_a \mathbf{6}_b + \mathbf{6}_b \mathbf{6}_a$ Strings stretched between the a^{th} and b^{th} stack lead to I_{ab} replicated chiral left-handed ⁴ fermions in the bifundamental representation $(N_a, \overline{N}_b)$. Additional light scalars may be present, with masses (12) in terms of angles determined by the wrapping numbers and the $\mathbf{T}^2$ moduli.

Generalization for compact spaces more general than the 6-torus will be discussed in section 3.2. We have therefore obtained a large class of four-dimensional theories with interesting non-abelian gauge symmetries and replicated charged chiral fermions. Hence compactifications with intersecting D6-branes provide a natural setup in which string theory can produce gauge sectors with the same rough features of the Standard Model. It is interesting to explore them further as possible phenomenological models, and construct explicit examples with spectrum as close as possible to the Standard Model.

3.1.2 Cancellation of RR tadpoles

String theories with open string sectors must satisfy a crucial consistency condition, known as cancellation of RR tadpoles. As mentioned above, D-branes act as sources for RR p -forms via the disk coupling $\int_{W_{p+1}} C_p$, see fig 9a. The consistency condition amounts to requiring the total RR charge of D-branes to vanish, as implied by Gauss law in a compact space (since RR field fluxlines cannot escape, figure 10). In our setup, the 3-cycle homology classes are

⁴Negative intersection numbers lead to a positive number of chiral fermions with right-handed chirality.




Figure 9: Disk diagrams couple D-branes to closed strings, and lead to tadpoles for RR fields.




Figure 10: In a compact space, fluxlines cannot escape and the total charge must vanish.

vectors of RR charges, hence the condition reads

$$[\Pi_{\text{tot}}] = \sum_a N_a [\Pi_a] = 0 \quad (15)$$

Equivalently, the condition of RR tadpole cancellation can be expressed as the requirement of consistency of the equations of motion for RR fields. In our situation, the terms of the spacetime action depending on the RR 7-form C_7 are

$$\begin{aligned} S_{C_7} &= \int_{\mathbf{M}_4 \times \mathbf{X}_6} H_8 \wedge *H_8 + \sum_a N_a \int_{\mathbf{M}_4 \times \Pi_a} C_7 = \\ &= \int_{\mathbf{M}_4 \times \mathbf{X}_6} C_7 \wedge dH_2 + \sum_a N_a \int_{\mathbf{M}_4 \times \mathbf{X}_6} C_7 \wedge \delta(\Pi_a) \end{aligned} \quad (16)$$

where H_8 is the 8-form field strength, H_2 its Hodge dual, and $\delta(\Pi_a)$ is a bump 3-form localized on Π_a in $\mathbf{X}_6$. The equations of motion read

$$dH_2 = \sum_a N_a \delta(\Pi_a) \quad (17)$$

The integrability condition is obtained by taking this equation in homology, yielding (15).

It is useful to describe the latter interpretation of RR cancellation condition from the viewpoint of the compactified theory. In this language, there exist components of the KK

reduction of C_7 leading to 4d RR 4-forms. Taking a basis of 3-cycles in $\mathbf{X}_6$, $[\Sigma_i]$, these are the zero modes $C_{4,i} = \int_{[\Sigma_i]} C_7$. Being 4d 4-forms, these fields do not have kinetic terms in their 4d action, and they only appear in linear tadpole terms of the form

$$S_{C_{4,i}} = [\Pi_a] \cdot [\Sigma_i] \int_{\mathbf{M}_4} C_{4,i} \quad (18)$$

The equations of motion then imply that the coefficient of the tadpole must vanish $[\Pi_a] \cdot [\Sigma_i] = 0$, namely it is not a condition on the field but a consistency condition for the model. For a complete basis of $[\Sigma_i]$, this implies (15).

3.1.3 Anomaly cancellation

Cancellation of RR tadpoles in the underlying string theory configuration implies cancellation of four-dimensional chiral anomalies in the effective field theory in our configurations. Recall that the chiral piece of the spectrum is given by I_{ab} chiral fermions in the representation $(N_a, \overline{N}_b)$ of the gauge group $\prod_a U(N_a)$.

Cubic non-abelian anomalies

The $SU(N_a)^3$ cubic anomaly is proportional to the number of fundamental minus anti-fundamental representations of $SU(N_a)$, hence it is proportional to $\sum_b I_{ab} N_b$.

It is easy to check this vanishes due to RR tadpole cancellation: Starting with (15), we consider the intersection of $[\Pi_{\text{tot}}]$ with any $[\Pi]$ to get

$$0 = [\Pi_a] \cdot \sum_b N_b [\Pi_b] = \sum_b N_b I_{ab} \quad (19)$$

as claimed.

It is interesting to notice that RR tadpole cancellation is slightly stronger than cancellation of cubic non-abelian anomalies. In fact, the former requires that the number of fundamental minus antifundamentals vanishes even for the cases $N_a = 1, 2$, where no gauge theory anomaly exists. This observation will turn out relevant in phenomenological model building.

Cancellation of mixed anomalies

The $U(1)_a$ - $SU(N_b)^2$ mixed anomalies also cancel as a consequence of RR tadpole cancellation. They do so in a trickier way, namely the anomaly receives two non-zero contributions which cancel each other ⁵, see fig 11.

⁵Mixed gravitational triangle anomalies cancel automatically, without Green-Schwarz contributions.




Figure 11: Triangle and Green-Schwarz diagrams contributing to the mixed $U(1)$ - non-abelian anomalies.

The familiar field theory triangle diagrams give a contribution which (after using RR tadpole conditions) is

$$A_{ab} \simeq N_a I_{ab} \quad (20)$$

On the other hand, the theory contains contributions from Green-Schwarz diagrams, where the gauge boson of $U(1)_a$ mixes with a 2-form which subsequently couples to two gauge bosons of $SU(N_b)$.

The coupling between $U(1)_a$ and a 2-form $\int_{[\Pi_a]} C_5$ arises from the coupling

$$N_a \int_{D6_a} C_5 \wedge \text{tr } F_a \xrightarrow{KK} N_a \int_{\mathbf{M}_4} B_a \wedge \text{tr } F_a \quad (21)$$

There are also scalars $\phi_b = \int_{[\Pi_b]} C_3$ which couple to $SU(N_b)$ gauge bosons via

$$\int_{D6_b} C_3 \wedge \text{tr } F_a^2 \xrightarrow{KK} \int_{\mathbf{M}_4} \phi_b \text{tr } F_b^2 \quad (22)$$

One can get the duality relation $dB_a = I_{ab} * d\phi_b$ and check that the contribution to the anomaly is proportional to

$$A'_{ab} \simeq -N_a I_{ab} \quad (23)$$

leading to a cancellation between both kinds of contributions.

An important observation is that any $U(1)$ gauge boson with $B \wedge F$ couplings gets massive, with mass roughly of the order of the string scale, see fig 12. Such $U(1)$'s disappear as gauge symmetries from the low-energy effective field theory, but remain as global symmetries, unbroken in perturbation theory.

$$\text{U(1)}_a \text{ --- } \text{U(1)}_a = m^2 A_u^2$$

Figure 12: The $B \wedge F$ couplings lead to a $U(1)$ gauge boson mass term.

3.1.4 A Standard Model - like example

Consider a configuration of D6-branes on $\mathbf{T}^6$ defined by the following wrapping numbers

$$\begin{aligned} N_1 &= 3 & (1, 2) & (1, -1) & (1, -2) \\ N_2 &= 2 & (1, 1) & (1, -2) & (-1, 5) \\ N_3 &= 1 & (1, 1) & (1, 0) & (-1, 5) \\ N_4 &= 1 & (1, 2) & (-1, 1) & (1, 1) \\ N_5 &= 1 & (1, 2) & (-1, 1) & (2, -7) \\ N_6 &= 1 & (1, 1) & (3, -4) & (1, -5) \end{aligned}$$

The intersection numbers are

$$\begin{aligned} I_{12} &= 3 & I_{13} &= -3 & I_{14} &= 0 & I_{15} &= 0 & I_{16} &= -3 \\ I_{23} &= 0 & I_{24} &= 6 & I_{25} &= 3 & I_{26} &= 0 & I_{34} &= -6 \\ I_{35} &= -3 & I_{36} &= 0 & I_{45} &= 0 & I_{46} &= 6 & I_{56} &= 3 \end{aligned}$$

A $U(1)$ linear combination, playing the role of hypercharge, remains massless

$$Q_Y = -\frac{1}{3} Q_1 - \frac{1}{2} Q_2 - Q_3 - Q_5 \quad (24)$$

The chiral fermion spectrum, with charges with respect to the Standard Model - like gauge group, is

$$\begin{aligned} &SU(3) \times SU(2) \times U(1)_Y \times \dots \\ &3(3, 2)_{1/6} + 3(\bar{3}, 1)_{-2/3} + 3(\bar{3}, 1)_{1/3} + 6(1, 2)_{-1/2} + \\ &+ 3(1, 2)_{1/2} + 6(1, 1)_1 + 3(1, 1)_{-1} + 9(1, 1)_0 \end{aligned} \quad (25)$$

It is a quite nice toy model, very close to the Standard Model spectrum, but with additional matter, in particular six extra $SU(2)$ doublets. The origin of the later can be understood from RR tadpole cancellation: As mentioned above, they require the number of fundamentals

to equal that of antifundamentals, even for $SU(2)$. Since, to obtain three left-handed quarks we have chiral fermions in $3(3, \bar{2})$, the full model must contain nine fields $(1, 2)$, three of which correspond to left handed leptons and six of them remaining as exotic additional matter. In section 4.3.1 we will see that more advanced models, including orientifold planes, can avoid this difficulty.

3.2 Generalization beyond torus

Clearly the above setup is not restricted to toroidal compactifications. Indeed one may take any compact 6-manifold as internal space, for instance a Calabi-Yau threefold, which would lead to $4d \mathcal{N} = 2$ supersymmetry in the closed string sector. In this situation we should pick a set of 3-cycles Π_a on which we wrap N_a D6-branes (for instance special lagrangian 3-cycles of $\mathbf{X}_6$ if we are interested in preserving supersymmetry), making sure they satisfy the RR tadpole cancellation condition $\sum_a N_a [\Pi_a] = 0$.

The final open string spectrum (for instance, in the case of supersymmetric wrapped D6-branes) arises in two kinds of sectors

6_a - 6_a Leads to $U(N_a)$ gauge bosons ($\mathcal{N} = 1$ vector multiplets in the supersymmetric case) and $b_1(\Pi_a)$ real adjoint scalars (chiral multiplets in susy case).

6_a - 6_b + 6_b - 6_a We obtain I_{ab} chiral fermions in the representation $(N_a, \bar{N}_b)$ (plus light scalars, massless in supersymmetry preserving intersections). Here $I_{ab} = [\Pi_a] \cdot [\Pi_b]$.

Notice that the chiral spectrum is obtained in terms of purely topological information of the configuration, as should be the case.

The phenomenology of non-toroidal models will be quite similar to that of toroidal compactifications with D-branes, see next subsection. Thus, the later are in any event good toy model for many features of general compactifications with intersecting branes. This is particularly interesting since it is relatively difficult to construct explicit configurations of intersecting D6-branes in Calabi-Yau models (although some explicit examples have been discussed in [8, 9]).

3.3 Phenomenological features

We now turn to a brief discussion of the phenomenological properties natural in this setup [2].




Figure 13: .

- As we soon discuss, most models constructed in the literature are non-supersymmetric. It is possible but notoriously more difficult to construct fully $\mathcal{N} = 1$ supersymmetric models. Therefore, unless alternative solutions to the hierarchy model are provided, the best proposal is to consider non-susy models to have a low string scale $M_s \simeq \text{TeV}$ to avoid hierarchy.

- The proton is stable in these models, since the $U(1)$ within the $U(3)$ color factor plays the role of baryon number, and is preserved as a global symmetry, exactly unbroken in perturbation theory. Non-perturbative effects breaking it arise from euclidean D2-branes wrapped on 3-cycles, and have the interpretation of spacetime gauge theory instantons, hence reproducing the non-perturbative breaking of baryon number in the Standard Model. This feature of proton stability is most welcome in models with low fundamental scale.

- These models do *not* have a natural gauge coupling unification, even at the string scale. Each gauge factor has a gauge coupling controlled by the volume of the wrapped 3-cycle

$$\frac{1}{g_{YM,a}^2} = \frac{M_S^3 V_{\Pi_a}}{g_s} \quad (26)$$

Gauge couplings are related to geometric volumes, hence their experimental values can be adjusted/reproduced in concrete models, rather than predicted by the general setup.

- There exists a geometric interpretation for the spontaneous electroweak symmetry breaking. In explicit models, the Higgs scalar multiplet arises from the light scalars at intersections (which are massless in susy cases, and massive or tachyonic in non-susy cases). In intersecting D-brane configurations, vevs for scalars at intersections parametrize the possibility of recombining two intersecting cycles into a single smooth one, as shown in figure 13. In the process, the gauge symmetry is reduced, corresponding to a Higgs mechanism in the effective field theory. See [6] for further discussion.

- There is a natural exponential hierarchy of the Yukawa couplings. Yukawa couplings




Figure 14: Geometric origin of the hierarchy of Yukawa couplings for different generations.

among the scalar Higgs and chiral fermions at intersections arise at tree level in the string coupling from open string worldsheet instantons; namely from string worldsheets spanning the triangle with vertices at the intersections and sides on the D-branes. Their value is roughly given by e^{-A} , with A the triangle area in string units⁶. Since different families are located at different intersections, their triangles have areas increasing linearly with the family index, leading to an exponential Yukawa hierarchy, see fig 14. See [18] for further analysis of yukawa couplings in explicit models.

4 Supersymmetry

In this section we discuss the supersymmetry or not of models of intersecting D6-branes, and the role of supersymmetry in phenomenological model building with these configurations.

The first observation we can make is that all models above constructed are non-supersymmetric. One simple way to see it is that we start with type IIA string theory compactified on $\mathbf{X}_6$, and introduce D6-branes. Since RR tadpole cancellation requires that the total RR charge vanishes, we are forced to introduce objects with opposite RR charges, in a sense branes and antibranes, a notoriously non-supersymmetric combination.

An equivalent derivation of the result is as follows: If we would succeed in constructing a supersymmetric configuration of D6-branes, the system as a whole would be a supersymmetric BPS state of type IIA on $\mathbf{X}_6$. Since for a BPS state the tension is proportional to the RR charge, and the latter vanishes due to RR tadpole cancellation, the tension of the state must vanish. The only D6-brane configuration with zero tension is having no D6-brane at

⁶This is a good approximation only for large triangles, otherwise further worldsheet instanton contributions as well as fluctuations should be considered.

all. Hence the only supersymmetric configuration would be just type IIA on $\mathbf{X}_6$, with no brane at all.

These arguments suggest a way out of the impasse. In order to obtain $\mathcal{N} = 1$ supersymmetric compactifications we need to introduce objects with negative tension and negative RR charge, and which preserve the same supersymmetry as the D6-branes. Such objects exist in string theory and are orientifold 6-planes, O6-planes. Introduction of these objects leads to an interesting extension of the configurations above constructed, and will be studied in section 4.2. In particular we will use them to construct supersymmetric compactifications with intersecting D6-branes.

4.1 Issues on non-supersymmetric models

Before doing that, it is interesting to consider diverse difficulties one encounters in non-supersymmetric models, and the extent to which they can be satisfactorily solved.

4.1.1 Tachyons at intersections

Even if each stack of D6-branes preserve some supersymmetry, the preserved susy may be different for different stacks. In these situations, the spectrum of open strings at intersections is generically non-supersymmetric, and some light scalar at the intersection may be tachyonic. Such tachyons signal an instability against recombination of the two sets of intersecting branes into a single smooth one. The initially considered configuration is not really stable.

There are two possible ways to face these tachyons. The first possibility is to consider configurations where all scalar fields at all intersections are non-tachyonic, leading to non-supersymmetric models which are nevertheless stable against small perturbations. These are at best local minima, since as argued above, there always exists a global supersymmetric minimum, which is the type IIA vacuum, and corresponds to having no D6-brane at all. The metastable vacua may tunnel (by nucleation of D8-brane bubbles, see footnote 8 in [19]) to this global minimum; they however provide reasonable enough models if sufficiently long-lived.

A second possibility is to employ tachyons of this kind, with the right quantum numbers, to trigger electroweak symmetry breaking. Although the string scale should be higher than the weak scale, the gauge symmetry breaking scale can be adjusted to around 100 GeV by fine-tuning the angles at the corresponding intersection. See [6] for further discussion of the

Higgs as a tachyon.

4.1.2 Hierarchy problem

Despite their simplicity for model building, non-supersymmetric models of intersecting D6-branes on $\mathbf{T}^6$ suffer a hierarchy problem. In order to avoid large corrections to the weak scale, it is natural to lower the string scale to around 1 TeV. The large value of the 4d Planck scale should therefore arise as a derived value due to some large internal dimension. However, if one attempts to increase any direction in $\mathbf{T}^6$, too light KK replicas of Standard Model gauge bosons are generated, because the corresponding D6-branes span all such directions. In other words, there is no direction in $\mathbf{T}^6$ which is transverse to all Standard Model D6-branes.

This is however a mild problem, in the sense that it is very specific to toroidal models. It is easy to imagine more general Calabi-Yau compactifications where the 3-cycles wrapped by the Standard Model branes are localized in a small region of the Calabi-Yau, and may be of small volume even for large Calabi-Yau volume. Some steps towards explicit realizations of this idea have been taken in [8, 9].

4.1.3 NSNS tadpoles

The most difficult issue in non-supersymmetric string vacua (even in general situations, not restricted to the case of intersecting D6-brane models) is the existence of uncanceled tadpoles for NSNS fields (graviton, dilaton, moduli).

In contrast with RR tadpoles, NSNS tadpoles do *not* signal an inconsistency of the theory. Reviewing the arguments in section 3.1.2, they differ in the fact that NSNS fields do have kinetic terms, hence the equations of motion can be satisfied by balancing the tadpole term with the kinetic term. The existence of a nonzero tadpole in a particular background means that it is not a correct background for the theory, and should be consequently corrected. The corrected backgrounds are usually curved geometries which must be computed in an α' expansion, resulting in quite complicated final models, sometimes (but not always [20]) involving naked singularities and strong coupling regions (see [21]).

In view of this situation two different attitudes have been taken. It is fair to say that none of them solves the issue of NSNS tadpoles in a fully satisfactory manner; after all, it is related to the difficult and unsolved questions of breaking of supersymmetry.

- Stick to $\mathcal{N} = 1$ supersymmetric models, to avoid NSNS tadpoles in the string construction. This approach obviously ‘ignores’ supersymmetry breaking in the real world, since

once this is implemented the issue of NSNS tadpoles would re-arise.

- Ignore the issue of NSNS tadpoles completely! This approach is justified by recognizing that we really do not know the complete equations of motion of string theory, hence it is possible that a background which does not solve the classical equations of motion still provides a good approximation to the solution to the full string equations of motion.

A good field theory analogy has been proposed by C. Bachas [11]. Consider a 4d $U(1)$ gauge field theory with charged fermions and a charged massless complex scalar with potential $V = |\phi|^4$. The solution to the classical equations of motion is $\phi = 0$, around which the theory has unbroken gauge symmetry and massless fermion and scalar excitations. A configuration not solving the equations of motion is $\phi = v \neq 0$, which sits on the slope of the scalar potential and therefore leads to a non-zero ϕ -tadpole. Ignoring the tadpole, the physics around this configuration is that of spontaneously broken gauge symmetry and massive fermions and scalar.

Interestingly enough, in many situations the second configuration provides the best approximation to the real physics of the system once quantum corrections are taken into account, since they may develop a mass term for ϕ , and a minimum of the corrected scalar potential with non-zero vev for ϕ . The complete physics of the system is better approximated by a configuration *not* solving the classical equations of motion. Analogously [11], *‘Perhaps the price for getting a good description of the low energy world from string theory may be to allow for small metric, dilaton and moduli tadpoles in the classical description of the groundstate’*.

Our purpose in the remaining is to discuss some of the most interesting models constructed within each approach. Before doing that, it will be convenient to introduce some background material on certain useful objects, the O6-planes.

4.2 Detour: Orientifold 6-planes

Consider type IIA theory on $\mathbf{X}_6$, and mod out the configuration by Ω (worldsheet orientation reversal) times a $\mathbf{Z}_2$ symmetry g of $\mathbf{X}_6$. The set of fixed points of g form an orientifold plane, namely a subspace of spacetime where the orientation of the string can flip.

A simple example, on which we center henceforth, is provided by $\mathbf{X}_6 = \mathbf{T}^6$ with g given by the action $y_i \rightarrow -y_i$, where y_i are the vertical direction on each $\mathbf{T}^2$. This is a symmetry for rectangular two-tori (see figure 15), or for two-tori tilted by a specific angle [3], see




Figure 15: Orientifold 6-planes in the orientifold quotient of IIA on $\mathbf{T}^6$ by Ωg , with $g : y_i \rightarrow -y_i$.




Figure 16: Cycles and their orientifold images in a rectangular and tilted 2-tori.

below. For figure 15, the set of fixed points is given by x_i arbitrary, $y_i = 0, R_{y_i}/2$, hence has 8 components. Since each has seven dimensions (counting also the $\mathbf{M}_4$ piece), they correspond to O6-planes wrapped on the 3-cycle with wrapping numbers $(n_i, m_i) = (1, 0)$.

One may consider introducing D6-branes in the above orientifold quotient of IIA on $\mathbf{T}^6$. The configurations are described as in the toroidal situation, by specifying the multiplicities N_a and wrapping numbers (n_a^i, m_a^i) of the D6-brane stacks, with two novelties:

First, in order to have a configuration invariant under Ωg we need to introduce orientifold images (denoted D6 $_{a'}$ -branes) of the D6 $_a$ -branes. They have multiplicity N_a and wrapping numbers $(n_a^i, -m_a^i)$ for rectangular 2-tori, or $(n_a^i, -n_a^i - m_a^i)$ for tilted tori, see fig 16. To simplify the latter case we introduce $\tilde{m}_a = m_a + n_a/2$, so that branes and images have wrapping numbers $(n_a, \tilde{m}_a)$ and $(n_a, -\tilde{m}_a)$ respectively.

Second, the O6-planes preserve the same supersymmetry as D6-branes on the same 3-cycle, and they carry negative tension and negative RR charge (-4 in D6-brane charge units). Consequently, they contribute to the RR tadpole condition, which now reads (for rectangular tori, i.e. 8 O6-planes)

$$\sum_a N_a [\Pi_a] + \sum_a N_a [\Pi_{a'}] - 4 \times 8 [\Pi_{O6}] = 0 \quad (27)$$

The open string spectrum in orientifolded models now contains new sectors. It is given by

aa+a'a' Contains $U(N_a)$ gauge bosons and superpartners

ab+ba+b'a'+a'b' Contains I_{ab} chiral fermions in the representation $(N_a, \overline{N}_b)$, plus light scalars.

ab'+b'a+ba'+a'b Contains $I_{ab'}$ chiral fermions in the representation (N_a, N_b) , plus light scalars.

aa'+a'a Contains certain numbers $n_{\square\square}, n_{\square\Box}$ of chiral fermions in the two-index symmetric resp. antisymmetric representations, with $n_{\square\square} + n_{\square\Box} = I_{aa'}$.

As expected, the new RR tadpole conditions in the presence of O6-planes guarantee the cancellation of 4d anomalies of the new chiral spectrum, in analogy with the toroidal case ⁷.

4.3 Advanced model building

4.3.1 Models in the non-susy setup

In this section we consider some of the phenomenologically most interesting examples of non-supersymmetric models, ‘ignoring’ the issue of the corresponding NSNS tadpoles. What follows is a brief sketch of the ideas in [4], to which we refer the reader for details.

Consider type IIA on $\mathbf{T}^6$ modded out by Ωg with $g : y_i \rightarrow -y_i$, and introduce four stacks of D6-branes (plus images) leading to a gauge group

$$U(3)_a \times U(2)_b \times U(1)_c \times U(1)_d \quad (28)$$

and with the only non-zero intersection numbers being

$$\begin{aligned} I_{ab} &= 1 & I_{ab'} &= 2 & I_{ac} &= -3 & I_{ac'} &= -3 \\ I_{bd} &= 0 & I_{bd'} &= -3 & I_{cd} &= -3 & I_{cd'} &= 3 \end{aligned}$$

One can check that the number of fundamentals equals that of antifundamentals even for $SU(2)$ and $U(1)$ factors. For instance, for $SU(2)$ we have left handed quarks in the representation $2(3, \overline{2}) + (\overline{1}, 2)$ and leptons in $3(1, 2)$. Notice that the model does not require six extra doubles, avoiding this feature of the purely toroidal model thanks to the existence of the new kind of bifundamental representation (N_a, N_b) .

⁷In the orientifold case, mixed gravitational anomalies may receive Green-Schwarz contributions [7]

In [4] a large class of explicit 3-cycles in $\mathbf{T}^6$ with those intersection numbers were constructed, and for which there is a unique massless $U(1)$ linear combination which plays the role of hypercharge. The resulting complete chiral spectrum is

$$\begin{aligned}
& SU(3)_c \times SU(2)_w \times U(1)_Y \\
3 \times & [(3, 2)_{1/6} + (\bar{3}, 1)_{1/3} + (\bar{3}, 1)_{-2/3} + \\
& + (1, 2)_{-1/2} + (1, 1)_1 + (1, 1)_0]
\end{aligned} \tag{29}$$

Namely *just* the Standard Model (plus right handed neutrinos).

Much of the recent activity in string model building has centered on the search of similarly successful models in non-toroidal geometries [8, 9], or supersymmetric realizations of these intersection numbers. In many respects, the search is on!

4.3.2 Supersymmetric models

In this section we review the first supersymmetric 4d chiral model of intersecting D6-branes, in [7] to which we refer the reader for details (see [22] for additional models).

In order to obtain supersymmetric models, one needs a sufficient number of O6-planes in the construction. One of the simplest possibilities is the Ωg orientifold of the $\mathbf{T}^6/(\mathbf{Z}_2 \times \mathbf{Z}_2)$ orbifold. The rules for model building are similar to the orientifolded tori case, with additional care to implement the $\mathbf{Z}_2$ orbifold actions.

Skipping the details, let us simply mention that the conditions to have all D6-brane intersections supersymmetric and cancel RR tadpoles are restrictive and do not allow for a systematic classification of models. Among the most interesting ones in [7], let us simply mention a configuration of six D6-brane stacks (plus Ωg images), with wrapping numbers

$$\begin{aligned}
N_1 = 8 & \quad (0, 1) \times (0, -1) \times (2, -1) & \rightarrow U(1) \\
N_2 = 2 & \quad (1, 0) \times (1, 0) \times (2, -1) \\
N_3 = 4 & \quad (1, 0) \times (1, -1) \times (1, 1) & \rightarrow U(2) \\
N_4 = 2 & \quad (1, 0) \times (0, 1) \times (0, -1) \\
N_5 = 6 + 2 & \quad (1, -1) \times (1, 0) \times (1, 0) & \rightarrow U(3) \times U(1) \\
N_6 = 4 & \quad (0, 1) \times (1, 0) \times (0, -1)
\end{aligned} \tag{30}$$

The configuration is $\mathcal{N} = 1$ supersymmetric for a suitable choice of the $\mathbf{T}^2$ radii ratios.

As shown above, the model contains a subgroup with the non-abelian structure of the Standard Model. It also contains one linear combination of the above $U(1)$'s which remains massless, and which plays the role of hypercharge. The final chiral spectrum, with respect to this Standard Model subgroup is

$$\begin{aligned}
& SU(3)_c \times SU(2)_w \times U(1)_Y \\
3 \times & [(3, 2)_{1/6} + (\bar{3}, 1)_{1/3} + (\bar{3}, 1)_{-2/3} + \\
& + (1, 2)_{-1/2} + (1, 1)_1 + (1, 1)_0]
\end{aligned} \tag{31}$$

plus *chiral* exotic matter. Chiral exotics render the model unrealistic, but the nice Standard Model - like subsector make it a good toy model of supersymmetric model building with intersecting D6-branes. Some of its phenomenological features have been described in [23].

5 Final comments

We have shown that compactifications with D6-branes wrapped on intersecting 3-cycles provide a large class of 4d models with non-abelian gauge symmetries and chiral fermions. A first lesson to be drawn is that D-branes in string theory *do* allow for phenomenologically realistic consistent theories realizing the brane world idea, and therefore providing an alternative to other scenarios, like weakly coupled heterotic theory or Horava-Witten theory compactifications.

We have briefly discussed some of the interesting phenomenological features of the present setup, like the possibility to lower the string scale, the stability of the proton, and the interesting hierarchy of yukawa couplings. Also, although we did not discuss them, the models (in particular the supersymmetric ones) have a rich structure of duality relations with other interesting string and M theory constructions, like M-theory on G_2 manifolds, or D-branes with worldvolume magnetic fluxes. Hopefully further research of these models and their properties will shed light on the difficult issues that remain to be addressed, most notably the issue of supersymmetry breaking and/or NSNS tadpoles.

Acknowledgements

I would like to thank G. Aldazabal, M. Cvetič, S. Franco, L. E. Ibáñez, R. Rabadán and G. Shiu for collaboration in these topics, and the organizers of the TMR meeting for creating an stimulating environment. I also thank M. González for her kind support and encouragement.

References

- [1] R. Blumenhagen, L. Goerlich, B. Kors, D. Lust, ‘Noncommutative compactifications of type I strings on tori with magnetic background flux’, JHEP 0010 (2000) 006, hep-th/0007024.
- [2] G. Aldazabal, S. Franco, L. E. Ibanez, R. Rabadan, A. M. Uranga, ‘D=4 chiral string compactifications from intersecting branes’, J. Math. Phys. 42 (2001) 3103, hep-th/0011073; ‘Intersecting brane worlds’, JHEP 0102 (2001) 047, hep-ph/0011132.
- [3] R. Blumenhagen, B. Kors, D. Lust, ‘Type I strings with F flux and B flux’, JHEP 0102 (2001) 030, hep-th/0012156.
- [4] L. E. Ibanez, F. Marchesano, R. Rabadan, ‘Getting just the standard model at intersecting branes’, JHEP 0111 (2001) 002, hep-th/0105155.
- [5] S. Forste, G. Honecker, R. Schreyer, ‘Orientifolds with branes at angles’, JHEP 0106 (2001) 004, hep-th/0105208;
R. Blumenhagen, B. Kors, D. Lust, T. Ott, ‘The standard model from stable intersecting brane world orbifolds’, Nucl. Phys. B616 (2001) 3, hep-th/0107138;
D. Cremades, L. E. Ibanez, F. Marchesano, ‘SUSY Quivers, Intersecting Branes and the Modest Hierarchy Problem’, hep-th/0201205;
C. Kokorelis, ‘GUT model hierarchies from intersecting branes’, hep-th/0203187; ‘New standard model vacua from intersecting branes’, hep-th/0205147; ‘Exact standard model compactifications from intersecting branes’, hep-th/0206108.
- [6] D. Cremades, L. E. Ibanez, F. Marchesano, ‘Intersecting brane models of particle physics and the Higgs mechanism’, JHEP 0207 (2002) 022, hep-th/0203160.
- [7] M. Cvetič, G. Shiu, A. M. Uranga, ‘Chiral four-dimensional N=1 supersymmetric type 2A orientifolds from intersecting D6 branes’, Nucl. Phys. B615 (2001) 3, hep-th/0107166; ‘Three family supersymmetric standard - like models from intersecting brane worlds’, Phys. Rev. Lett. 87 (2001) 201801, hep-th/0107143.
- [8] R. Blumenhagen, V. Braun, B. Körs, D. Lüst, ‘Orientifolds of K3 and Calabi-Yau manifolds with intersecting D-branes’, JHEP 0207 (2002) 026, hep-th/0206038.
- [9] A. M. Uranga, ‘Local models for intersecting brane worlds’, hep-th/0208014.

- [10] R. Blumenhagen, L. Gorlich, B. Kors, ‘Supersymmetric 4-D orientifolds of type IIA with D6-branes at angles’, JHEP 0001 (2000) 040, hep-th/9912204;
S. Forste, G. Honecker, R. Schreyer, ‘Supersymmetric $Z(N) \times Z(M)$ orientifolds in 4-D with D branes at angles’, Nucl. Phys. B593 (2001) 127, hep-th/0008250.
- [11] C. Bachas, ‘A Way to break supersymmetry’, hep-th/9503030.
- [12] C. Angelantonj, I. Antoniadis, E. Dudas, A. Sagnotti, ‘Type I strings on magnetized orbifolds and brane transmutation’, Phys. Lett. B489 (2000) 223, hep-th/0007090.
- [13] J. Polchinski, ‘Dirichlet Branes and Ramond-Ramond charges’, Phys. Rev. Lett. 75 (1995) 4724, hep-th/9510017.
- [14] N. Arkani-Hamed, S. Dimopoulos, G.R. Dvali ‘The hierarchy problem and new dimensions at a millimeter’, Phys.Lett.B429:263-272,1998 , hep-ph/9803315; I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos, G.R. Dvali ‘New dimensions at a millimeter to a fermi and superstrings at a TeV’, Phys.Lett.B436:257-263,1998, hep-ph/9804398.
- [15] M. R. Douglas, G. W. Moore, ‘D-branes, quivers, and ALE instantons’, hep-th/9603167;
M. R. Douglas, B. R. Greene, D. R. Morrison, ‘Orbifold resolution by D-branes’, Nucl. Phys. B506 (1997) 84, hep-th/9704151.
- [16] G. Aldazabal, L. E. Ibáñez, F. Quevedo, A. M. Uranga, ‘D-branes at singularities: A Bottom up approach to the string embedding of the standard model’, JHEP 0008 (2000) 002, hep-th/0005067;
D. Berenstein, V. Jejjala, R. G. Leigh, ‘The Standard model on a D-brane’, Phys. Rev. Lett. 88 (2002) 071602, hep-ph/0105042;
L. F. Alday, G. Aldazabal, ‘In quest of just the standard model on D-branes at a singularity’, JHEP 0205 (2002) 022, hep-th/0203129;
- [17] M. Berkooz, M. R. Douglas, R. G. Leigh, ‘Branes intersecting at angles’, Nucl. Phys. B480 (1996) 265, hep-th/9606139.
- [18] D. Cremades, L.E. Ibanez, F. Marchesano, ‘Towards a theory of quark masses, mixings and CP violation’, hep-ph/0212064.
- [19] A. M. Uranga, ‘Localized instabilities at conifolds’, hep-th/0204079.

- [20] E. Dudas, J. Mourad, C. Timirgaziu, ‘Time and space dependent backgrounds from nonsupersymmetric strings’, hep-th/0209176.
- [21] E. Dudas, J. Mourad, ‘Brane solutions in strings with broken supersymmetry and dilaton tadpoles’, Phys. Lett. B486 (2000) 172, hep-th/0004165; R. Blumenhagen, A. Font, ‘Dilaton tadpoles, warped geometries and large extra dimensions for nonsupersymmetric strings’, Nucl. Phys. B599 (2001) 241, hep-th/0011269.
- [22] R. Blumenhagen, L. Gorlich, T. Ott, ‘Supersymmetric intersecting branes on the type 2A T6 / Z(4) orientifold’, hep-th/0211059;
M. Cvetič, I. Papadimitriou, G. Shiu, ‘Supersymmetric three family SU(5) grand unified models from type IIA orientifolds with intersecting D6-branes’, hep-th/0212177.
- [23] M. Cvetič, P. Langacker, G. Shiu, ‘Phenomenology of a three family standard like string model’, Phys. Rev. D66 (2002) 066004, hep-ph/0205252; ‘A Three family standard - like orientifold model: Yukawa couplings and hierarchy’, Nucl. Phys. B642 (2002) 139, hep-th/0206115.